

Sorption Reversibility of Cadmium from Aqueous Solutions on Natural Firoozkoh Zeolite

F Pooladi¹ and M Hamidrpour^{1*}

(Received: July 8-2013 ; Accepted : Nov. 16-2013)

Abstract

Removal of boron from aqueous environments (soil and water) is difficult, because it is present as $B(OH)_3$ and $B(OH)_4^-$ species. This research was done to study the sorption of boron by HDTMA-modified zeolite. The sorption of B on modified zeolite was studied as a function of pH (B concentration: 1 and 10 mg L⁻¹) in the range of 6-9.5, and as a function of ionic strength (0.03 and 0.06 M Ca(NO₃)₂ or Mg(NO₃)₂) at a constant B concentration of 5 mg L⁻¹. Sorption isotherm was performed for the solutions containing initial B concentration in the range of 1-15 mg L⁻¹ using a 24h batch equilibration experiment. The results revealed that surfactant-modified zeolite exhibited the best performance at pH 9.5, and sorption of B increased with the increase of suspension pH. Greater B adsorption in the Ca system over the Mg system was clearly observed for the modified zeolite. Sorption isotherm of B were well described by the Freundlich and Langmuir models but the Freundlich sorption model described the interaction between B and the mineral material better than the Langmuir model. Maximum sorption capacity (q_{max}) of the sorbent was 120 mmol kg⁻¹. The experimental data showed that HDTMA-modified zeolite used in this study had a reasonable sorption capacity for B.

Keywords: Boron, Sorption isotherm, Modified zeolite, Removal.

1. Dept. of Soil Sci., Isf. Univ. of Technol., Isfahan, Iran.

2. Dept. of Soil Sci., Vali-e-Asr University of Rafsanjan, Rafsanjan, Iran.

*: Corresponding Author, Email: mohsen_hamidpour@yahoo.com