

Analysis of 2,4-D Concentration in Silty Clay Soil Under Two Irrigation Regimes Using LEACHP and PRZM-3 Models

M. Noshadi*, S. Jamshidi and F. Foroharfar¹

(Received: Nov. 16-2013 ; Accepted : Oct. 04-2015)

Abstract

Evaluation of pollutant transportation in soil is important from different environmental aspects such as soil and groundwater contamination. The purpose of this study is to measure 2, 4-D concentrations in a silty loam soil under two different treatments (normal and deficit irrigation) in a corn field and simulate the results using the PRZM-3 and LEACHP models. Total concentrations of 2, 4-D in the soil profile in 8, 13, 23, 30, 37, and 57 days after application for normal irrigation were 18.5, 16.36, 11.67, 10.47, 8.47 and 3.2 mg kg⁻¹, respectively. For these dates, PRZM-3 model simulated 18.5, 16.36, 11.67, 10.47, 8.47 and 3.2 mg kg⁻¹ of 2,4-D, respectively and LEACHP model simulated 23.34, 20.93, 16.7, 16.3, 12.9 and 11.41 mg kg⁻¹ of 2, 4-D, respectively. Total concentrations of 2, 4-D in the mentioned dates for deficit irrigation were 20.2, 16.7, 11.22, 10.05, 8.8 and 7.3 mg kg⁻¹, respectively. For these dates, PRZM-3 model simulation results were 21.9, 19.89, 14.2, 10.62, 9.6 and 8.22 mg kg⁻¹, respectively and LEACHP model simulation results were 25.22, 21.3, 19.43, 18.58, 18 and 16.27 mg kg⁻¹, respectively. The simulation results showed that performance of PRZM-3 model was better than LEACHM model in both treatments. In this research, the half-lives of 2, 4-D for 0-10 cm and 10-20 cm of soil depth were 7 and 33 days in a normal irrigation, and 9 and 34.65 days in a deficit irrigation, respectively.

Keywords: Simulation, Normal irrigation, Deficit irrigation.

1. Dept. of Water Eng., Shiraz Univ., Shiraz, Iran.

*: Corresponding Author, Email: noshadi@shirazu.ac.ir