

Effect of Endophytic Fungus, *Piriformospora indica*, on Barley Resistance to Lead

F. Karimi¹, M. Sepehri², M. Afuni³ and M. A. Hajabbasi⁴

(Received: Nov. 27-2011 ; Accepted : Oct. 21-2014)

Abstract

By modifying plants at genetical, physiological and ecological levels, entophytic fungi as the most important soil microorganisms have a pronounced growth-promoting activity and also increase plant resistance to biotic and abiotic stresses. This research was undertaken to evaluate the potential of *P. indica* to increase barley (*Hordeum vulgare* L.) resistance to lead (pb). Therefore, a greenhouse experiment with two fungus treatments (non-inoculated and *P. indica* inoculated) and five levels of pb (0, 25, 50, 100 and 500 mg/kg) with three replications was conducted based on a factorial design. Measurement of shoot and root dry weight showed that the growth of *P. indica*-colonized plants at all levels of pb treatments was higher ($P < 0.05$) than that of the corresponding controls. Also, chlorophyll concentration of inoculated plants with *P. indica* was superior to non-inoculated plants. In addition, the results showed that in contrast to the plant shoot, lead concentration in the root of *P. indica*-colonized plants was higher than the non-inoculated controls.

Keyword: *Piriformospora indica*, Lead, Barley.

1. Dept. of Soil Sci., College of Agric., Isf. Univ. Technol., Isfahan, Iran.

*: Corresponding Author, Email: fatemeh.karimi@rocketmail.com