Simulation of Fluoride Transport in a Calcareous Soil Using HYDRUS-1D

E. Chavoshi^{1*}, M. Afyuni² and M. A. Hajabbasi²

(Received: July 23-2014; Accepted: Feb. 21-2015)

Abstract

Transport of fluoride and consumption of groundwater with excess fluoride concentrations poses a health threat to millions of people around the world. The objective of this study was to simulate transport of fluoride (F) using HYDRUS-1D model. The study was conducted in lysimeters at Lavark research station site in Isfahan. The treatments consisted of two concentrations of F (157 and 315 mg kg⁻¹). The duration of the study was 125 days. Some of soil physical and chemical properties, soluble F and total F concentration were determined during the study. The results showed the transport of F in calcareous soil profiles. This may be due to the high pH and desorption of F ion as a result of repulsion by the more negatively charged soil surfaces. The highest concentration of total F and water soluble F were observed in the 10 cm surface soil layer. The concentration of F decreased with increased soil depth. The correlation coefficient was significant between the water soluble fluoride and the total fluoride (1% level). Also, the difference between the observed t- value and a critical value on the t distribution is statistically insignificant. It showed that the model simulated successfully water soluble F concentration in the soil profile.

Keywords: Fluoride, model, HYDRUS-1D, calcareous soil.

^{1.} Dept. of Soil Sci., Collage of Agric., Isfahan (Khorasgan) Branch, Islamic Azad Univ., Isfahan, Iran.

^{2.} Dept. of Soil Sci., College of Agric., Isf., Univ. of Technol., Isfahan, Iran.

^{*:} Corresponding Author, Email: chavoshie@yahoo.com